arXiv:1606.02805v1 [gr-qc] 9 Jun 2016

Axisymmetric Stationary Spacetimes of Constant Scalar Curvature in Four Dimensions

Rosikhuna F. Assafari[♯,a)], Bobby E. Gunara[§,♯,b)] [1], Hasanuddin[‡,c)], and Abednego Wiliardy [♯,d)]

[§] *Indonesian Center for Theoretical and Mathematical Physics (ICTMP)*
and
[♯] *Theoretical Physics Laboratory*
Theoretical High Energy Physics and Instrumentation Research Group,
Faculty of Mathematics and Natural Sciences,
Institut Teknologi Bandung
Jl. Ganesha no. 10 Bandung, Indonesia, 40132
and
[‡] *Department of Physics,*
Faculty of Mathematics and Natural Science,
Tanjungpura University,
Jl. Prof. Dr. H. Hadari Nawawi, Pontianak, Indonesia, 78124
[a)]rosikhuna@yahoo.co.id, [b)]bobby@fi.itb.ac.id, [c)]hasanuddin@physics.untan.ac.id,
[d)]abednego.wiliardy@gmail.com

Abstract

In this paper we construct a special class of four dimensional axissymmetric stationary spacetimes whose Ricci scalar is constant in the Boyer-Lindquist coordinates. The first step is to construct Einstein metric by solving a modified Ernst equation for nonzero cosmological constant. Then, we modify the previous result by adding two additional functions to the metric to obtain a more general metric of constant scalar curvature which are not Einstein.

1 INTRODUCTION

The aim of this paper is to construct a special class of four dimensional axissymmetric stationary spacetimes of constant scalar curvature in the Boyer-Lindquist coordinates. First, we discuss the construction of Einstein spacetimes (or known to be Kerr-Einstein spacetimes) with nonzero cosmological constant by solving a modified Ernst equation for nonzero cosmological constant. In other words, we re-derive the Carter's result in [1]. Then, we proceed to construct the spaces of constant scalar curvature which are not Einstein by modifying the previous result, namely, we add two additional functions to the metric to have a more

[1] Corresponding author.

general form of the metric but it has the structure of polynomial, namely, it is a quartic polynomial.

The structure of the paper can be mentioned as follows. In section 2 we give a quick review on axissymmetric stationary spacetimes. Then, we begin our construction of Kerr-Einstein spacetimes in section 3. We discuss the construction of axissymmetric stationary spacetimes of constant scalar curvature in section 4.

2 Axissymmetric Stationary Spacetimes: A Quick Review

Suppose we have a metric in the general form

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu \ , \tag{1}$$

defined on a four dimensional spacetimes $\mathbf{M^4}$, where x^μ parametrizes a local chart on $\mathbf{M^4}$ and $\mu, \nu = 0, ..., 3$. Then, we simplify the case as follows. In a stationary axisymmetric spacetime, the time coordinate t and the azimuthal angle φ are considered to be x^0 and x^1 respectively. A stationary axisymmetric metric is invariant under simultaneous transformations $t \to -t$ and $\varphi \to -\varphi$ which yields

$$g_{02} = g_{03} = g_{12} = g_{13} = 0 \ , \tag{2}$$

and moreover, all non-zero metric components depend only on $x^2 \equiv r$ and $x^3 \equiv \theta$. The latter condition implies $g_{23} = 0$ and the metric (1) can be simplified into [2, 3]

$$ds^2 = -e^{2\nu}dt^2 + e^{2\psi}(d\varphi - \omega dt)^2 + e^{2\mu_2}dr^2 + e^{2\mu_3}d\theta^2 \ , \tag{3}$$

where $(\nu, \psi, \omega, \mu_2, \mu_3) \equiv (\nu(r,\theta), \psi(r,\theta), \omega(r,\theta), \mu_2(r,\theta), \mu_3(r,\theta))$. In the following we list the non-zero components of Christoffel symbol related to the metric (3):

$$\Gamma^0{}_{02} = \nu_{,2} - \frac{1}{2}\omega\omega_{,2}\ e^{2(\psi-\nu)} \ , \ \Gamma^0{}_{03} = \nu_{,3} - \frac{1}{2}\omega\omega_{,3}e^{2(\psi-\nu)} \ ,$$

$$\Gamma^0{}_{12} = \frac{1}{2}\omega_{,2}e^{2(\psi-\nu)} \ , \ \Gamma^0{}_{13} = \frac{1}{2}\omega_{,3}e^{2(\psi-\nu)} \ ,$$

$$\Gamma^1{}_{20} = -\omega\left(\psi_{,2} - \nu_{,2}\right) - \frac{1}{2}\omega_{,2}(1 + \omega^2 e^{2(\psi-\nu)}) \ ,$$

$$\Gamma^1{}_{12} = \psi_{,2} + \frac{1}{2}\omega\omega_{,2}\ e^{2(\psi-v)} \ , \ \Gamma^1{}_{13} = \psi_{,3} + \frac{1}{2}\omega\omega_{,3}\ e^{2(\psi-\nu)} \ ,$$

$$\Gamma^2{}_{00} = \nu_{,2}\ e^{2(v-\mu_2)} - \omega(\omega_{,2} + \omega\psi_{,2})e^{2(\psi-\mu_2)} \ , \tag{4}$$

$$\Gamma^2{}_{01} = \left(\frac{1}{2}\omega_{,2} + \omega\psi_{,2}\right)e^{2(\psi-\mu_2)} \ , \ \Gamma^2{}_{11} = -\psi_{,2}e^{2(\psi-\mu_2)} \ ,$$

$$\Gamma^2{}_{22} = \mu_{2,2} \ , \ \Gamma^2{}_{23} = \mu_{2,3} \ , \ \Gamma^2{}_{33} = -\mu_{3,2} \ e^{2(\mu_2-\mu_3)} \ ,$$

$$\Gamma^3{}_{00} = \nu_{,3} e^{2(\nu-\mu_3)} - \omega(\omega_{,3} + \omega\psi_{,3}) e^{2(\psi-\mu_3)} \ ,$$

$$\Gamma^3{}_{01} = \left(\frac{1}{2}\omega_{,3} + \omega\psi_{,3}\right) e^{2(\psi-\mu_3)} \ , \ \Gamma^3{}_{11} = -\psi_{,3} \ e^{2(\psi-\mu_3)} \ ,$$

$$\Gamma^3{}_{22} = -\mu_{2,3} \ e^{2(\mu_2-\mu_3)} \ , \ \Gamma^3{}_{23} = \mu_{3,2} \ , \ \Gamma^3{}_{33} = \mu_{3,3} \ ,$$

and the non-zero components of Ricci tensor:

$$\begin{aligned}
R_{00} &= e^{2(\nu-\mu_2)} \left(\nu_{,2,2} + v_{,2}(\psi + \nu - \mu_2 + \mu_3)_{,2} - \frac{1}{2}\omega^2{}_{,2} e^{2(\psi-\nu)} \right) \\
&\quad + e^{2(\nu-\mu_3)} \left(\nu_{,3,3} + \nu_{,3}(\psi + \nu + \mu_2 - \mu_3)_{,3} - \frac{1}{2}\omega^2{}_{,3} e^{2(\psi-\nu)} \right) \\
&\quad - \omega e^{2(\psi-\mu_2)} \left(\omega_{,2,2} + \omega_{,2}(3\psi - \nu - \mu_2 + \mu_3)_{,2} \right) \\
&\quad - \omega e^{2(\psi-\mu_3)} \left(\omega_{,3,3} + \omega_{,3}(3\psi - \nu + \mu_2 - \mu_3)_{,3} \right) \\
&\quad - \omega^2 e^{2(\psi-\mu_2)} \left(\psi_{,2,2} + \psi_{,2}(\psi + \nu - \mu_2 + \mu_3)_{,2} + \frac{1}{2}\omega^2{}_{,2} e^{2(\psi-\nu)} \right) \\
&\quad - \omega^2 e^{2(\psi-\mu_3)} \left(\psi_{,3,3} + \psi_{,3}(\psi + \nu + \mu_2 - \mu_3)_{,3} + \frac{1}{2}\omega^2{}_{,3} e^{2(\psi-\nu)} \right) \ , \\
R_{01} &= \frac{1}{2} e^{2(\psi-\mu_2)} \left(\omega_{,2,2} + \omega_{,2}(3\psi - \nu - \mu_2 + \mu_3)_{,2} \right) \\
&\quad + \frac{1}{2} e^{2(\psi-\mu_3)} \left(\omega_{,3,3} + \omega_{,3}(3\psi - \nu + \mu_2 - \mu_3)_{,3} \right) \\
&\quad + \omega e^{2(\psi-\mu_2)} \left(\psi_{,2,2} + \psi_{,2}(\psi + \nu - \mu_2 + \mu_3)_{,2} + \frac{1}{2}\omega^2{}_{,2} e^{2(\psi-\nu)} \right) \\
&\quad + \omega e^{2(\psi-\mu_3)} \left(\psi_{,3,3} + \psi_{,3}(\psi + \nu + \mu_2 - \mu_3)_{,3} + \frac{1}{2}\omega^2{}_{,3} e^{2(\psi-\nu)} \right) \ , \quad (5) \\
R_{11} &= - e^{2(\psi-\mu_2)} \left(\psi_{,2,2} + \psi_{,2}(\psi + \nu - \mu_2 + \mu_3)_{,2} + \frac{1}{2}\omega^2{}_{,2} e^{2(\psi-\nu)} \right) \\
&\quad - e^{2(\psi-\mu_3)} \left(\psi_{,3,3} + \psi_{,3}(\psi + \nu + \mu_2 - \mu_3)_{,3} + \frac{1}{2}\omega^2{}_{,3} e^{2(\psi-\nu)} \right) \ , \\
R_{22} &= - (\psi_{,2,2} + \psi_{,2}(\psi - \mu_2)_{,2} - (\nu_{,2,2} + v_{,2}(\nu - \mu_2)_{,2} \\
&\quad - e^{2(\mu_2-\mu_3)} \left(\mu_{2,3,3} + \mu_{2,3}(\psi + \nu + \mu_2 - \mu_3)_{,3} \right) \\
&\quad - \left(\mu_{3,2,2} + \mu_{3,2}(\mu_3 - \mu_2)_{,2} \right) + \frac{1}{2}\omega^2{}_{,2} e^{2(\psi-v)} \ , \\
R_{23} &= - (\psi_{,2,3} + \psi_{,2}(\psi - \mu_2)_{,3} - (\nu_{,2,3} + \nu_{,2}(v - \mu_2)_{,3} + \mu_{3,2}(\psi - \nu)_{,3} \\
&\quad + \frac{1}{2}\omega_{,2}\omega_{,3} e^{2(\psi-\nu)} \ , \\
R_{33} &= - (\psi_{,3,3} + \psi_{,3}(\psi - \mu_3)_{,3} - (\nu_{,3,3} + \nu_{,3}(v - \mu_3)_{,3} \\
&\quad - e^{2(\mu_3-\mu_2)} \left(\mu_{3,2,2} + \mu_{3,2}(\psi + \nu - \mu_2 + \mu_3)_{,3} \right) \\
&\quad - \left(\mu_{2,3,3} + \mu_{2,3}(\mu_2 - \mu_3)_{,3} \right) + \frac{1}{2}\omega^2{}_{,3} e^{2(\psi-\nu)} \ .
\end{aligned}$$

Then, Ricci scalar can be obtained as

$$
\begin{aligned}
-R &= 2e^{-2\mu_2}\Big(\psi_{,2,2}+\psi_{,2}(\psi-\mu_2+\mu_3)_{,2}+\psi_{,2}\nu_{,2}+\nu_{,2,2}+\nu_{,2}(\nu-\mu_2+\mu_3)_{,2} \\
&\quad +\mu_{3,2,2}+\mu_{3,2}(\mu_3-\mu_2)_{,2}-\frac{1}{4}{\omega^2}_{,2}e^{2(\psi-\nu)}\Big) \\
&\quad +2e^{-2\mu_3}\Big(\psi_{,3,3}+\psi_{,3}(\psi+\mu_2-\mu_3)_{,3}+\psi_{,3}\nu_{,3}+\nu_{,3,3}+\nu_{,3}(\nu+\mu_2-\mu_3)_{,3} \\
&\quad +\mu_{2,3,3}+\mu_{2,3}(\mu_2-\mu_3)_{,3}-\frac{1}{4}{\omega^2}_{,3}e^{2(\psi-\nu)}\Big)\ , \qquad (6)
\end{aligned}
$$

where we have defined

$$
f_{,\mu}\equiv\frac{\partial f}{\partial x^\mu}\ ,\quad f_{,\mu,\nu}\equiv\frac{\partial^2 f}{\partial x^\mu\partial x^\nu}\ . \qquad (7)
$$

3 EINSTEIN SPACETIMES

In this section, we construct a class of axissymmetric spacetimes satisfying Einstein condition

$$
R_{\mu\nu}=\Lambda g_{\mu\nu}\ , \qquad (8)
$$

with Λ is named cosmological constant, yielding the following coupled nonlinear equations:

$$
\left(e^{3\psi-\nu-\mu_2+\mu_3}\omega_{,2}\right)_{,2}+\left(e^{3\psi-\nu+\mu_2-\mu_3}\omega_{,3}\right)_{,3}=0\ , \qquad (9)
$$

$$
(\psi+\nu)_{,2,3}-(\psi+\nu)_{,2}\mu_{2,3}-(\psi+\nu)_{,3}\mu_{3,2}+\psi_{,2}\psi_{,3}+\nu_{,2}\nu_{,3}=\frac{1}{2}e^{2(\psi-\nu)}\omega_{,2}\ \omega_{,3}\ , \qquad (10)
$$

$$
\left(e^{\mu_3-\mu_2}\left(e^\beta\right)_{,2}\right)_{,2}+\left(e^{\mu_2-\mu_3}\left(e^\beta\right)_{,3}\right)_{,3}=-2\Lambda e^{\beta+\mu_2+\mu_3}\ , \qquad (11)
$$

$$
\left(e^{\beta-\mu_2+\mu_3}(\psi-\nu)_{,2}\right)_{,2}+\left(e^{\beta+\mu_2-\mu_3}(\psi-\nu)_{,3}\right)_{,3}=-e^{3\psi-\nu}\left(e^{\mu_3-\mu_2}{\omega_{,2}}^2+e^{\mu_2-\mu_3}{\omega_{,3}}^2\right), \qquad (12)
$$

$$
\begin{aligned}
4e^{\mu_3-\mu_2}\left(\beta_{,2}\mu_{3,2}+\psi_{,2}\nu_{,2}\right)-4e^{\mu_2-\mu_3}\left(\beta_{,3}\mu_{2,3}+\psi_{,3}\nu_{,3}\right) &= 2e^{-\beta}\left[\left(e^{\mu_3-\mu_2}\left(e^\beta\right)_{,2}\right)_{,2}+\left(e^{\mu_2-\mu_3}\left(e^\beta\right)_{,3}\right)_{,3}\right] \\
&\quad -e^{2(\psi-\nu)}\left(e^{\mu_3-\mu_2}{\omega_{,2}}^2-e^{\mu_2-\mu_3}{\omega_{,3}}^2\right)\ , \qquad (13)
\end{aligned}
$$

where we have defined

$$
\beta\equiv\psi+\nu\ . \qquad (14)
$$

This class of solutions is called Kerr-(anti) de Sitter solutions.

3.1 The Functions μ_2 and μ_3

First of all, we simply take e^{μ_2} as

$$
e^{\mu_2}=\frac{(r^2+a^2\cos^2\theta)^{\frac{1}{2}}}{{\Delta_r^{(0)}}^{\frac{1}{2}}}\ , \qquad (15)
$$

where $\Delta_r^{(0)} \equiv \Delta_r^{(0)}(r)$ and a is a constant related to the angular momentum of a black hole [3]. Next, we assume that the function $e^{2(\mu_3-\mu_2)}$ and $e^{2\beta}$ are separable as

$$\begin{aligned} e^{2(\mu_3-\mu_2)} &= \Delta_r^{(0)} \frac{\sin^2\theta}{\Delta_\theta^{(0)}} \ , \\ e^{2\beta} &= \Delta_r^{(0)} \Delta_\theta^{(0)} \ , \end{aligned} \tag{16}$$

with $\Delta_\theta^{(0)} \equiv \Delta_\theta^{(0)}(\theta)$. Thus, (11) can be cast into the form

$$\left[\Delta_r^{(0)\frac{1}{2}}\left(\Delta_r^{(0)\frac{1}{2}}\right)_{,2}\right]_{,2} + \frac{1}{\sin\theta}\left[\frac{\Delta_\theta^{(0)\frac{1}{2}}}{\sin\theta}\left(\Delta_\theta^{(0)\frac{1}{2}}\right)_{,3}\right]_{,3} = -2\Lambda\left(r^2 + a^2\cos^2\theta\right). \tag{17}$$

Employing the variable separation method, we then obtain

$$\begin{aligned} \Delta_r^{(0)} &= -\frac{\Lambda}{3}r^4 + c_1 r^2 + c_2 r + c_3 \ , \\ \Delta_\theta^{(0)} &= -\frac{\Lambda}{3}a^2\cos^4\theta - c_1\cos^2\theta - c_4\cos\theta + c_5 \ , \end{aligned} \tag{18}$$

where c_i $i = 1, ..., 5$, are real constant. To make a contact with [1], one has to set c_i to be

$$\begin{aligned} c_1 &= -\frac{\Lambda}{3}a^2 \ , \ c_2 = -2M \ , \ c_3 = a^2 \ , \\ c_4 &= 0 \ , \ c_5 = 1 \ , \end{aligned} \tag{19}$$

such that we have

$$\begin{aligned} \Delta_r^{(0)} &= -\frac{\Lambda}{3}r^2\left(r^2+a^2\right) + r^2 - 2Mr + a^2 \ , \\ \Delta_\theta^{(0)} &= \left(1 + \frac{\Lambda}{3}a^2\cos^2\theta\right)\sin^2\theta \ . \end{aligned} \tag{20}$$

3.2 The Functions (ω, ν, ψ) and Ernst Equation

To obtain the explicit form of (ω, ν, ψ), we have to transform (9) and (12) into a so called Ernst equation with nonzero Λ using (20). This can be structured as follows.

First, we introduce a pair of functions (Φ, Ψ) via

$$\begin{aligned} \Phi_{,2} &= e^{2(\psi-\nu)}\Delta_\theta^{(0)}\omega_{,p} \ , \\ \Phi_{,p} &= -e^{2(\psi-\nu)}\Delta_r^{(0)}\omega_{,2} \ , \\ \Psi &\equiv e^{\psi-\nu}\Delta_r^{(0)\frac{1}{2}}\Delta_\theta^{(0)\frac{1}{2}} \ , \end{aligned} \tag{21}$$

where $p \equiv \cos\theta$. Then, (9) and (12) can be cast into

$$\begin{aligned}
\Psi\left[(\Delta_r^{(0)}\Phi_{,2})_{,2} + (\Delta_\theta^{(0)}\Phi_{,p})_{,p}\right] &= 2\Delta_r^{(0)}\Psi_{,2}\Phi_{,2} + 2\Delta_\theta^{(0)}\Psi_{,p}\Phi_{,p} \ , \qquad (22)\\
\Psi\left[(\Delta_r^{(0)}\Psi_{,2})_{,2} + (\Delta_\theta^{(0)}\Psi_{,p})_{,p}\right] &= \Delta_r^{(0)}\left[(\Psi_{,2})^2 - (\Phi_{,2})^2\right] - \Delta_\theta^{(0)}\left[(\Psi_{,p})^2 - (\Phi_{,p})^2\right] \ ,
\end{aligned}$$

respectively. Defining a complex function $Z \equiv \Psi + \mathrm{i}\Phi$, (22) can be rewritten in Ernst form

$$ReZ\left[(\Delta_r^{(0)}Z_{,2})_{,2} + (\Delta_\theta^{(0)}Z_{,p})_{,p}\right] = \Delta_r^{(0)}(Z_{,2})^2 + \Delta_\theta^{(0)}(Z_{,p})^2 \ . \qquad (23)$$

Note that one could obtain another solution of (23), say $\tilde{Z} \equiv \tilde{\Psi} + \mathrm{i}\tilde{\Phi}$ by a conjugate transformation

$$\begin{aligned}
\tilde{\Psi} &= \frac{\Delta_r^{(0)\frac{1}{2}}\Delta_\theta^{(0)\frac{1}{2}}}{\tilde{\chi}} \ ,\\
\tilde{\Phi}_{,2} &= \frac{\Delta_\theta^{(0)}}{\tilde{\chi}^2}\tilde{\omega}_{,3} \ , \qquad (24)\\
\tilde{\Phi}_{,3} &= -\frac{\Delta_r^{(0)}}{\tilde{\chi}^2}\tilde{\omega}_{,2} \ ,
\end{aligned}$$

where

$$\begin{aligned}
\tilde{\chi} &\equiv \frac{e^{\nu-\psi}}{e^{2(\nu-\psi)} - \omega^2} \ ,\\
\tilde{\omega} &\equiv \frac{\omega}{e^{2(\nu-\psi)} - \omega^2} \ . \qquad (25)
\end{aligned}$$

In the latter basis, we find

$$\begin{aligned}
\tilde{\Psi} &= \frac{\Delta_r^{(0)} - a^2\Delta_\theta^{(0)}}{r^2 + a^2\cos^2\theta} \ ,\\
\tilde{\Phi} &= \frac{2aM\cos\theta}{r^2 + a^2\cos^2\theta} + \frac{2\Lambda}{3}ar\cos\theta \ . \qquad (26)
\end{aligned}$$

After some computation, we conclude that [1]

$$\begin{aligned}
e^{2\psi} &= \frac{\left(r^2+a^2\right)^2\Delta_\theta^{(0)} - \Delta_r^{(0)}a^2\sin^4\theta}{r^2 + a^2\cos^2\theta} \ ,\\
e^{2\nu} &= \frac{(r^2 + a^2\cos^2\theta)\Delta_\theta^{(0)}\Delta_r^{(0)}}{\left(r^2+a^2\right)^2\Delta_\theta^{(0)} - \Delta_r^{(0)}a^2\sin^4\theta} \ , \qquad (27)\\
\omega &= \frac{a(r^2+a^2)\Delta_\theta^{(0)} - a\,\sin^2\theta)\Delta_r^{(0)}}{\left(r^2+a^2\right)^2\Delta_\theta^{(0)} - \Delta_r^{(0)}a^2\sin^4\theta} \ .
\end{aligned}$$

4 SPACETIMES OF CONSTANT RICCI SCALAR

In this section we extend the previous results to the case of spaces of constant Ricci scalar, namely

$$R = g^{\mu\nu} R_{\mu\nu} = k \ , \tag{28}$$

where k is a constant. To have an explicit solution, we simply replace $\Delta_r^{(0)}$ and $\Delta_\theta^{(0)}$ in (16), (20), and (27) by

$$\begin{aligned} \Delta_r &= \Delta_r^{(0)} + f(r) \ , \\ \Delta_\theta &= \Delta_\theta^{(0)} + h(\theta) \ . \end{aligned} \tag{29}$$

Then, inserting these modified functions mentioned above to (28), we simply have

$$-k(r^2 + a^2\cos^2\theta) = (\Delta_r)_{,2,2} + \frac{1}{\sin\theta}\left[\frac{(\Delta_\theta)_{,3}}{\sin\theta}\right]_{,3} \ , \tag{30}$$

which gives

$$-(k - 4\Lambda)(r^2 + a^2\cos^2\theta) = 2\left[f^{\frac{1}{2}}\left(f^{\frac{1}{2}}\right)_{,2}\right]_{,2} + \frac{2}{\sin\theta}\left[\frac{h^{\frac{1}{2}}}{\sin\theta}\left(h^{\frac{1}{2}}\right)_{,3}\right]_{,3} . \tag{31}$$

The solution of (31) is given by

$$\begin{aligned} f(r) &= -\frac{1}{12}(k - 4\Lambda) r^4 + \frac{1}{2}C_1 r^2 + C_2 r + C_3 \ , \\ h(\theta) &= -\frac{1}{12}(k - 4\Lambda) a^2\cos^4\theta \ - \frac{1}{2}C_1\cos^2\theta \ - C_4\cos\theta \ + C_5 \ , \end{aligned} \tag{32}$$

where $C_i, \ i = 1, ..., 5$, are real constant. It is worth mentioning some remarks as follows. First, the functions $f(r)$ and $g(\theta)$ have the same structure as in (18), namely they are quartic polynomials with respect to r and $\cos\theta$, respectively. Second, the constant Λ here is no longer the cosmological constant. Finally, Einstein spacetimes can be obtained by setting $k = 4\Lambda$ and $C_3 = a^2 C_5$ with other C_i i=1, 2, 4, are free constants.

Now we can state our main result as follows.

Theorem 1 *Suppose we have an axissymmetric spacetime* $\mathbf{M^4}$ *endowed with metric*

$$ds^2 = e^{2\nu}dt^2 - e^{2\psi}(d\varphi - \omega dt)^2 - e^{2\mu_2}dr^2 - e^{2\mu_3}d\theta^2 \ , \tag{33}$$

satisfying

$$
\begin{aligned}
e^{2(\mu_3-\mu_2)} &= \Delta_r \frac{\sin^2\theta}{\Delta_\theta} \ , \\
e^{2\beta} &= \Delta_r \Delta_\theta \ , \\
e^{2\psi} &= \frac{\left(r^2+a^2\right)^2 \Delta_\theta - \Delta_r a^2 \sin^4\theta}{r^2+a^2\cos^2\theta} \ , \\
e^{2\nu} &= \frac{(r^2+a^2\cos^2\theta)\Delta_\theta \Delta_r}{\left(r^2+a^2\right)^2 \Delta_\theta - \Delta_r a^2 \sin^4\theta} \ , \\
\omega &= \frac{a(r^2+a^2)\Delta_\theta - a\,\sin^2\theta \Delta_r}{\left(r^2+a^2\right)^2 \Delta_\theta - \Delta_r a^2 \sin^4\theta} \ ,
\end{aligned}
\tag{34}
$$

where Δ_r and Δ_θ are given by (29). Then, there exist a familiy of spacetimes of constant scalar curvature with

$$
\begin{aligned}
\Delta_r &= -\frac{1}{3}\Lambda r^2 a^2 + r^2 - 2Mr + a^2 - \frac{k}{12} r^4 + \frac{1}{2} C_1 r^2 + C_2 r + C_3 \ , \qquad (35) \\
\Delta_\theta &= -\cos^2\theta \ + \frac{\Lambda}{3} a^2 \cos^2\theta \ - \frac{k}{12} a^2 \cos^4\theta \ - \frac{1}{2} C_1 \cos^2\theta \ - C_4 \cos\theta \ + C_5 + 1 \ ,
\end{aligned}
$$

where $C_i,\ i = 1, ..., 5$, are real constant. The metric (33) becomes Einstein if $k = 4\Lambda$ and $C_3 = a^2 C_5$ for all C_i.

The norm of Riemann tensor for the case at hand in general has the form

$$
\begin{aligned}
R^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta} &= \frac{384 r^4}{(r^2+a^2\cos^2\theta)^6} \Bigg(2a^2 C_4 \cos\theta \Big(-a^2 C_5 + r(C_2 - 2M) + C_3 \Big) \\
&\quad + \Big(-a^2 C_5 + r(aC_4 + C_2 - 2M) + C_3 \Big) \Big(-C_5 a^2 + r(-aC_4 + C_2 - 2M) + C_3 \Big) \Bigg) \\
&\quad + \frac{192 r^2}{(r^2+a^2\cos^2\theta)^5} \Bigg(a^2 C_4 \cos\theta \left(3a^2 C_5 - 4C_2 r - 3C_3 + 8Mr \right) \\
&\quad - 5r(C_2 - 2M) \left(C_3 - a^2 C_5 \right) - 2 \left(C_3 - a^2 C_5 \right)^2 \\
&\quad - 3r^2 (-aC_4 + C_2 - 2M)(aC_4 + C_2 - 2M) \Bigg) \\
&\quad + \frac{8}{(r^2+a^2\cos^2\theta)^4} \Bigg(6a^2 C_4 \cos\theta \Big(-a^2 C_5 + 3r(C_2 - 2M) + C_3 \Big) \\
&\quad + 30r(C_2 - 2M) \left(C_3 - a^2 C_5 \right) + 7 \left(C_3 - a^2 C_5 \right)^2 \\
&\quad + 27 r^2 (aC_4 + C_2 - 2M)(-aC_4 + C_2 - 2M) \Bigg) \\
&\quad - \frac{12}{(r^2+a^2\cos^2\theta)^3} (-aC_4 + C_2 - 2M)(aC_4 + C_2 - 2M) + \frac{k^2}{6}
\end{aligned}
\tag{36}
$$

which might have a negative value as observed in [4] for Kerr-Newman metric with $k = 0$. This is so because the spacetime metric is indefinite. The norm (36) shows that the spacetime has a real ring singularity at $r = 0$ and $\theta = \pi/2$ with radius a.

Generally, the metric described in Theorem 1 may not be related to Einstein general relativity since our method described above does not use the notion of energy-momentum tensor. To make a contact with general relativity, we could simply set some constants, for example, namely

$$C_1 = C_2 = C_4 = C_5 = 0 \ , \ C_3 = q^2 + g^2 \ , \tag{37}$$

with $k = 4\Lambda$ where q and g are electric and magnetic charges, respectively. This setup gives the Kerr-Newman-Einstein metric describing a dyonic rotating black hole with nonzero cosmological constant [5].

A SPACETIME CONVENTION

In this section we collect some spacetime quantities which are useful for the analysis in the paper.

Christoffel symbol:

$$\Gamma^{\lambda}{}_{\mu\nu} = \frac{1}{2} g^{\rho\sigma} \left(\partial_\mu g_{\rho\nu} + \partial_\nu g_{\rho\mu} - \partial_\rho g_{\mu\nu} \right) \ . \tag{38}$$

Riemann curvature tensor:

$$- R^{\rho}_{\mu\nu\sigma} = \partial_\sigma \Gamma^{\rho}{}_{\mu\nu} - \partial_\nu \Gamma^{\rho}{}_{\mu\sigma} + \Gamma^{\lambda}{}_{\mu\nu} \Gamma^{\rho}{}_{\lambda\sigma} - \Gamma^{\lambda}{}_{\mu\sigma} \Gamma^{\rho}{}_{\lambda\nu} \ . \tag{39}$$

Ricci tensor:

$$R_{\mu\nu} = R^{\rho}_{\mu\rho\upsilon} = \partial_\rho \Gamma^{\rho}{}_{\mu\nu} - \partial_\nu \Gamma^{\rho}{}_{\mu\rho} + \Gamma^{\lambda}{}_{\mu\nu} \Gamma^{\rho}{}_{\lambda\rho} - \Gamma^{\lambda}{}_{\mu\rho} \Gamma^{\rho}{}_{\lambda\nu} \ . \tag{40}$$

Ricci scalar:

$$R = g^{\mu\nu} R_{\mu\nu} \ . \tag{41}$$

Acknowledgments

The work in this paper is supported by Riset KK ITB 2014-2016 and Riset Desentralisasi DIKTI-ITB 2014-2016.

References

[1] B. Carter, "Black holes equilibrium states," in "Proceedings, Ecole d'Et de Physique Thorique: Les Astres Occlus : Les Houches, ed. by B. DeWitt, C. M. DeWitt, (Gordon and Breach, New York, 1973).

[2] S. Chandrasekhar, "The Kerr Metric and Stationary Axis-Symmetric Gravitational Field," Proc. Roy. Soc. Lond. A **358**, 405 (1978).

[3] For a review see for example:
S. Chandrasekhar, "The mathematical theory of black holes," OXFORD, UK: CLARENDON (1985) 646 p and references therein.

[4] R. C. Henry, "Kretschmann scalar for a Kerr-Newman black hole," Astrophys. J. **535**, 350 (2000) [astro-ph/9912320].

[5] M. R. Setare and M. B. Altaie, "The Cardy-Verlinde formula and entropy of topological Kerr-Newman black holes in de Sitter spaces," Eur. Phys. J. C **30**, 273 (2003) [hep-th/0304072].